

Solvent Effects on Protein Motion and Protein Effects on Solvent Motion

Douglas J. Tobias¹, Mounir Tarek², Joseph E. Curtis³, I-Feng Kuo¹, Ali Razmara¹

¹Department of Chemistry, University of California, Irvine, California 92697-2025 USA

²Equipe de dynamique des assemblages membranaires, Unit Mixte de Recherche CNRS/UHP, Universite Henri Poincare-Nancy, B. P. 239, 54506 Vandoeuvre-les-Nancy Cedex, France

³NIST Center for Neutron Research, National Institute of Standards and Technology, Gaithersburg, Maryland 20899-8562 USA

Abstract

Proteins present rough, chemically heterogeneous, and dynamic surfaces to their solvent, which for most proteins is primarily aqueous[1]. Roughly a monolayer of water, the so-called hydration water is perturbed structurally and dynamically by interaction with the protein[2]. We will present results from several experimental and molecular dynamics simulation studies[2, 3, 4, 5] that illustrate the anomalous properties of hydration water, and demonstrate that hydration water is crucial for protein function. We will also discuss the coupling of functionally important protein motions to water dynamics[6], as well as how this coupling can be tuned by changing the solvent composition[7].

Acknowledgements:

This work is supported by the National Science Foundation (grant MCB-0078278).

References

- [1] Tobias, D. J.; Kuo, W. I.-F.; Razmara, A; Tarek, M. *In Water in Confining Geometries*, Devlin, J. P; Buch, V. Eds.; Springer: Amsterdam (2003).
- [2] Makarov, V.; Pettitt, B. M.; Feig, M.; *Accts. Chem. Res.*, 2002, **35**, 376, and references therein.
- [3] Settles, M; Doster, W.; *Faraday Disc.*, 1996, **103**, 269.
- [4] Fitter, J.; Lechner, R. E.; Dencher, N. A.; *J. Phys. Chem.*, 1999, **103**, 8036.
- [5] Tarek, M.; Tobias, D. J.; *Biophys. J.*, 2000, **79**, 3244.
- [6] Tarek, M.; Tobias, D. J.; *Phys. Rev. Lett.*, 2002, **88**, 138101.
- [7] Tsai, A. M.; Neumann, D. A.; Bell, L. N.; *Biophys. J.*, 2000, **79**, 2728.